

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

A-2

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ :C07C 229/26, 271/22, 309/73, C07D
211/60

A1

(11) International Publication Number:

WO 96/11181

(43) International Publication Date:

18 April 1996 (18.04.96)

(21) International Application Number: PCT/GB95/02385

(22) International Filing Date: 9 October 1995 (09.10.95)

(30) Priority Data:

9420243.9

7 October 1994 (07.10.94)

GB

9504928.4

10 March 1995 (10.03.95)

GB

(71) Applicant (for all designated States except US): CHIRO-
SCIENCE LIMITED [GB/GB]; Cambridge Science Park,
Milton Road, Cambridge CB4 4WE (GB).

(72) Inventor; and


(75) Inventor/Applicant (for US only): HUTTON, Gordon, Eric
[GB/GB]; Chiroscience Limited, Cambridge Science Park,
Milton Road, Cambridge CB4 4WE (GB).(74) Agent: GILL JENNINGS & EVERY; Broadgate House, 7
Eldon Street, London EC2M 7LH (GB).(81) Designated States: AM, AT, AU, BB, BG, BR, BY, CA, CH,
CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE,
KG, KP, KR, KZ, LK, LR, LT, LU, LV, MD, MG, MN,
MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK,
TJ, TM, TT, UA, UG, US, UZ, VN, European patent (AT,
BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL,
PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN,
ML, MR, NE, SN, TD, TG), ARIPO patent (KE, MW, SD,
SZ, UG).

Published

With international search report.

(54) Title: THE MANUFACTURE OF LEVOBUPIVACAINE AND ANALOGUES THEREOF FROM L-LYSINE

(57) Abstract

A novel compound of formula (X) is prepared from L-lysine, and
converted to Levobupivacaine by cyclisation.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LN	Lithuania	SN	Senegal
CN	China	LK	Sri Lanka	TD	Chad
CS	Czechoslovakia	LU	Luxembourg	TG	Togo
CZ	Czech Republic	LV	Latvia	TJ	Tajikistan
DE	Germany	MC	Monaco	TT	Trinidad and Tobago
DK	Denmark	MD	Republic of Moldova	UA	Ukraine
ES	Spain	MG	Madagascar	US	United States of America
FI	Finland	ML	Mali	UZ	Uzbekistan
FR	France	MN	Mongolia	VN	Viet Nam
GA	Gabon				

**THE MANUFACTURE OF LEVOBUPIVACAINE AND ANALOGUES
THEREOF FROM L-LYSINE**

Field of the Invention

5 This invention relates to a cost-effective process for the conversion, via diazotisation methodology, of L-lysine to a carboxanilide precursor of levobupivacaine and related compounds.

Background of the Invention

10 Compounds of formula 1 wherein R is a methyl, n-propyl or n-butyl group are widely used as local anaesthetics. Biological studies have shown that the (S)-enantiomers of such compounds (e.g. levobupivacaine wherein R is n-butyl) display lower cardiotoxicity than the corresponding racemates whilst maintaining the same anaesthetic potency,
15 and are therefore more beneficial for clinical uses. Thus there is a requirement for efficient processes to manufacture compounds of formula 1 in the form of single enantiomers. The process embodied by the present invention employs a chirality pool approach to levobupivacaine,
20 commencing from L-lysine, an inexpensive starting material which is readily available in bulk.

Although L-lysine has been converted through to optically enriched L-pipecolic acid and esters thereof by diazotisation and cyclisation reactions, cyclisation of an
25 intermediate to form a piperidine-2-carboxanilide directly has hitherto not been reported. Furthermore, in the context of the present invention, these existing methods are hampered by an excessive number of steps required for manipulation of protecting groups in order to obviate
30 Walden inversion at the carboxyl-bearing centre, which leads to the formation of D-pipecolic acid.

An additional benefit of the invention is the provision of unnatural amino acids of formula 2 with (S) configuration, which are important as pharmaceutical
35 intermediates, e.g. for incorporation into physiologically active synthetic peptides. Compound 2a is commonly prepared by a multistep synthesis of the racemate

commencing from dihydropyran followed by enzymatic resolution of an ester derivative (e.g. *p*-nitrobenzyl). This process is inefficient with respect to the number of steps required and by the fact that resolution produces up to 50% of an unwanted enantiomer. Diazotisation of L-lysine and derivatives thereof has been identified as an alternative strategy for the preparation of 2-amino-6-hydroxyhexanoic acid and derivatives thereof. However, such an approach has hitherto only been applied to the synthesis of compound 2c, but this process, employing sodium nitroprusside, is low yielding (28%).

Copper complexation of lysine and similar amino-acids has been extensively used as a technique for temporary protection of the α -amino acid moiety, as disclosed by Ledger et al, Anstr. J. Chem. 18:933-5 (1965). This procedure facilitates selective transformation of the second amino group present in the side-chain, e.g. the attachment of a covalently-bonded amino-protecting group.

Summary of the Invention

This invention provides a cost-effective process for preparing compounds of formula 1, wherein R is H or alkyl, from the inexpensive starting material L-lysine, using a chirality pool approach, and involving scission of the terminal C-N bond of L-lysine. In particular, the present invention provides a practical and economical process for the manufacture of levobupivacaine from L-lysine, which is exemplified in Scheme 1. This process involves the preparation of novel compounds of formula (X) as defined in claim 1; either R' is a protecting group, in which case R is H in the product, which can then be alkylated as desired, or R' is alkyl and corresponds to R as alkyl.

Description of the Invention

Formula (X), and other formulae used herein, refer to compounds that have at least predominantly the given stereochemistry. They are at least substantially free of their optical antipode, e.g. in at least 50%, more preferably at least 70% or 99%, enantiomeric excess.

According to the particular reactions shown in Scheme 1, the α -N-benzyloxycarbonyl derivative of L-lysine is initially (a) subjected to diazotisation in the presence of acetic acid to afford (S)-2-(benzyloxycarbonyl-amino)-6-acetoxihexanoic acid (2b). Alternatively, Z may be any other blocking group, many examples of which are well known to the skilled man.

The next three steps of Scheme 1 proceed by way of novel compounds 3-5: (b) Condensation with 2,6-dimethylaniline gives compound 3; (c) cleavage of the O-acetyl group by methanolysis gives compound 4; and (d) tosylation gives compound 5. The OTs group shown in Scheme 1 is merely one example of many leaving groups which can be used and which are known to the skilled man. Examples are given in claim 2.

Finally, compound 5 is (a) subjected to hydrogenation in the presence of base to effect direct conversion to enantiomerically pure (S)-2',6'-dimethylpiperidine-2-carboxanilide 6. Variants of this process include conversion of the α -N-benzyloxycarbonyl (or other blocked) derivative of L-lysine to compound 6 via alkyl halide derivatives of formula 7. The conversion of compound 6 to levobupivacaine or other N-alkylated analogues can be carried out by methodology known to those skilled in the art, or as described in another Patent Application filed today, claiming priority from British Patent Application No. 9421478.0. Preferably, R is methyl, n-propyl, cyclopropyl or, most preferably, n-butyl.

A preferred aspect of the process summarised in Scheme 1 is the use of either sodium nitrite/sodium acetate/acetic acid or isoamyl nitrite/acetic acid as efficient reagent systems for the conversion of α -N-benzyloxycarbonyl or otherwise blocked L-lysine to the compound 2b, by formation of a diazonium intermediate followed by solvolysis of this species or of the carbocation formed by loss of N_2 . This discovery provides significant benefit over the prior art

process since the yields are much higher (60-87% of 2b compared to 28% 2c).

As a further feature of the invention, we have discovered that L-lysine can be cleanly and efficiently converted to (S)-2-amino-6-hydroxyhexanoic acid 2a in a process (Scheme 2) involving sequential formation of bis(lysinato)copper, diazotisation-solvolysis to generate the novel copper(II) complex 8, and cation-exchange to effect decomplexation. Diazotisation-solvolysis in such processes can be effected in aqueous media using either sodium nitroprusside at pH >10 or sodium nitrite, e.g. at pH <4.

More generally, NH_2 may be converted to any leaving group Y, e.g. halide, usually under anhydrous conditions, using known methodology. Depending on the stability of the metal complex, the reaction proceeds in one or more steps, to give a compound of the formula



in optically-enriched form. The free amine may be blocked at this stage, as a reactant for use in Scheme 1.

Such processes are advantageous over conventional diazotisations of L-lysine, which give mixtures of products arising from unselective transformation of either one, and in some cases both, of the two amine groups present.

The following Examples illustrate the invention. Examples 1 to 5 illustrate respective steps in Scheme 1; Example 6 illustrates Scheme 2.

Example 1

A stirred solution of N^{α} -benzyloxycarbonyl L-lysine (6.33 g, 22.6 mmol) in acetic acid (150 ml) was treated with sodium acetate (1.85 g) and then sodium nitrite (1.56 g x 3) at 40°C. The reaction was stirred for 3 hours at 40°C, cooled and the bulk of the acetic acid was removed in vacuo and the residues partitioned between water (100 ml) and dichloromethane (100 ml x 3). The combined organic extracts were dried (MgSO_4) and evaporated in vacuo to give compound 2b a yellow oil (6.35 g, 87%).

Example 2

Isoamyl nitrite (6.1 ml) was added to a stirred solution of N^α-benzyloxycarbonyl L-lysine (6.34 g, 22.6 mmol) in glacial acetic acid (35 ml) at 22°C. The mixture was heated at 50°C for 12 hours, then volatile material removed by distillation in vacuo (last traces removed via formation of azeotropic mixtures with toluene) to give compound 2b (4.38 g, 60%).

Example 3

A solution of compound 2b (6.35g, 19.7 mmol) in dichloromethane (150 ml) was treated sequentially with 2,6-dimethylaniline and a solution of dicyclohexylcarbodiimide (4.89 g) in dichloromethane (10 ml) at 18°C. The reaction was stirred for 24 hours. The precipitate which formed was filtered and washed with dichloromethane (100 ml). The organics were concentrated in vacuo to give compound 3 as waxy solid. Without further treatment, this material was dissolved in MeOH (150 ml) and solid K₂CO₃ (9.57 g) was added. The reaction was stirred for 24 hours then filtered and evaporated to leave a waxy solid which was chromatographed on silica gel with 1.5:1 EtOAc:heptane to give compound 4 as a colourless solid (2.22 g, 28%).

Example 4


p-Toluenesulphonyl chloride (0.4 g) was added to a solution of compound 4 (0.60 g, 1.7 mmol) and pyridine (1.4 ml) in dichloromethane (5 ml) and stirred for 4 hours. Dilute hydrochloric acid (1 N; 10 ml) was added, and after stirring for 18 hours the mixture was extracted with dichloromethane (50 ml). The organic solution was washed with aqueous NaOH (1 N; 10 ml), dried (MgSO₄) and evaporated in vacuo to give a yellow oil which was chromatographed on silica gel eluting with 1:1 EtOAc:heptane. This gave compound 5 as a colourless oil (0.76 g, 81%) which slowly solidifies as the tosyl derivative.

Example 5


A mixture of compound 5 (0.76 g, 1.4 mmol), Pd/C (0.076 g) and K_2CO_3 (0.47 g) in EtOH (15 ml) was placed under an atmosphere of hydrogen (balloon) and stirred vigorously for 3 hours. The reaction was filtered and the solids thoroughly washed with EtOH (5 ml). The combined filtrate and washings were concentrated in vacuo and the residue partitioned between dichloromethane (3 x 25 ml) and aqueous NaOH (1 N; 25 ml). The combined organic extracts were dried ($MgSO_4$) and concentrated in vacuo to give (S)-2',6'-dimethylpiperidine-2-carboxanilide 6 as a colourless solid (0.284g, 87%), which was shown by chiral HPLC analysis to have an optical purity of >98%.

Example 6

A suspension of L-lysine (0.931 g, 5.1 mmol) and $CuCO_3 \cdot Cu(OH)_2$ (1.24 g, 5.6 mmol) in water (15 ml) was heated under reflux for 5 minutes then cooled and filtered. The pH of the filtrate was adjusted to 4 with 2M H_2SO_4 . A solution of sodium nitrite (0.70 g, 10 mmol) in water (5 ml) was added dropwise over 10 minutes and the mixture was stirred at 24°C for 6 hours. Chelex 100 resin (40 g) was added to effect decomplexation. The resulting suspension was then filtered to give a solution of L-lysine (60%) and (S)-2-amino-6-hydroxyhexanoic acid 2 (30%).




(1)




(2)

2a: R=R'=H
 2b: R=Ac; R'=Z
 2c: R=H, R'=Z


SCHEME 1



SCHEME 2

CLAIMS

1. A carboxanilide represented by formula X



10 wherein R' is a removable amino-protecting group or alkyl, and Y is a leaving group.

2. A compound according to claim 1, wherein R' is an amino-protecting group.

15 3. A compound according to claim 1 or claim 2, wherein Y is N_2^+ , OH, O-acyl, -alkylsulphonyl, O-arylsulphonyl or halide.

4. A compound according to any preceding claim, wherein R' is benzyloxycarbonyl.

20 5. A compound according to claim 4, wherein Y is acetoxy.

6. A process for the preparation of (S)-2',6'-dimethylpiperidine-2-carboxanilide from L-lysine, which comprises the steps of conversion of L-lysine to a carboxanilide according to any of claims 1 to 5, and cyclisation to form a piperidine ring.

25 7. A process for the preparation of a carboxanilide according to any of claims 1 to 5, which comprises diazotisation-solvolysis of L-lysine.

8. A process according to claim 7, which comprises using sodium nitrite and sodium acetate in acetic acid.

30 9. A process according to claim 7, which comprises using isoamyl nitrite in acetic acid.

10. Use of a carboxanilide according to any of claims 1 to 5, or the product of a process according to any of claims 6 to 9, for the manufacture of levobupivacaine.

35

11. Use of a carboxanilide according to any of claims 1 to 5, or the product of a process according to any of claims 6 to 9, for the manufacture of ropivacaine.
12. A copper or other metal complex of (S)-2-amino-6-(Y-substituted)hexanoic acid, wherein Y is OH or a leaving group such as a halogen atom.
13. An aqueous solution of the complex according to claim 12.
14. A process for preparing (S)-2-amino-6-(Y-substituted)hexanoic acid, wherein Y is as defined in claim 12, which comprises converting L-lysine to bis(lysinato)copper or another metal complex, conversion of the terminal NH₂ group(s) to Y, e.g. by diazotisation-solvolysis, and cation-exchange to effect decomplexation.
15. A process according to claim 14, wherein diazotisation-solvolysis is effected using sodium nitroprusside at basic pH, preferably above 10.
16. A process according to claim 15, wherein diazotisation-solvolysis is effected using sodium nitrite at acidic pH, preferably below 5.

INTERNATIONAL SEARCH REPORT

ational Application No

PCT/GB 95/02385

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C07C229/26 C07C271/22 C07C309/73 C07D211/60

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07C C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	SYNTHESIS., 1982 STUTTGART DE, page 163 L. KISFALUDY ET AL. 'ONE-STEP SYNTHESIS OF L-PIPERIDINE-2-CARBOXYLIC ACID' cited in the application see the whole document ---	1-11
A	AUSTRALIAN JOURNAL OF CHEMISTRY, vol. 18, 1965 MELBOURNE, pages 933-935, R. KEDGER ET AL. 'THE USE OF SEQUESTERING AGENTS IN THE PREPARATION OF EPSILON-L-LYSINE AND DELTA-L-ORNITHINE DERIVATIVES' cited in the application see the whole document -----	12-16

☐ Further documents are listed in the continuation of box C.☐ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- *&* document member of the same patent family

Date of the actual completion of the international search

11 December 1995

Date of mailing of the international search report

21.12.95

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+ 31-70) 340-3016

Authorized officer

Sánchez García, J.M.

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☒ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☒ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☒ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.